



HD 49798: First contracting WD

S.Popov, S. Mereghetti, S. Blinnikov, A. Kuranov, L. Yungelson

MNRAS (2018, in press) arXiv: 1711.02449



HD 49798




alpha(2000) = 06h 48m 04.64s , delta(2000) = -44d 18' 59.3"
V = 8.30, B-V = -0.29, Spectral type: O6
constellation Puppis
about 650 parsecs from Earth

HD 49798 was discovered in 1964
a rare hydrogen-deficient O class subdwarf

$$\dot{M}_W = 3 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$$



X-ray source RX J0648.0-4418
Porb=1.55 days
Pspin=13.2 s
Mx=1.28+/-0.05 Msolar



System parameters


Parameter	Value	Units
Right Ascension	$6^h 48^m 4.7^s$	J2000
Declination	$-44^\circ 18' 58.4''$	J2000
Orbital period	1.547666(2)	d
$A_X \sin i$	9.79(19)	light-s
T^*	43961.243(15)	MJD
ν_0	0.0758480846(1)	Hz
$\dot{\nu}$	$1.24(2) \times 10^{-17}$	Hz s^{-1}
P_0	13.18424856(2)	s
\dot{P}	$-2.15(5) \times 10^{-15}$	s s^{-1}
T_0	48937.7681361	MJD
M_X	1.28(5)	M_\odot
M_C	1.50(5)	M_\odot

Mereghetti et al. (2016) 1603.01505



Mereghetti et al. (2011) 1105.6227

Problems with interpretation: a WD or a NS?



Before Pdot measurements all systems parameters have been well explained in the model of accreting WD.

Pdot measurements pointed to a NS instead of a WD, because with the observed accretion luminosity it is difficult to explain such significant Pdot of a WD.


$$j = \frac{2\pi\dot{v}IGM}{L_X R},$$

$$j_{WD} = 2.2 \times 10^{19} \left(\frac{L_X}{2 \times 10^{32} \text{ erg s}^{-1}} \right)^{-1} \text{ cm}^2 \text{ s}^{-1},$$

$$j_{NS} = 5.5 \times 10^{16} \left(\frac{L_X}{2 \times 10^{32} \text{ erg s}^{-1}} \right)^{-1} \text{ cm}^2 \text{ s}^{-1}.$$

$$\dot{M} > 2 \times 10^{16} \text{ g s}^{-1}, \quad \text{in the case of a WD,}$$

$$\dot{M} > 2 \times 10^{11} \text{ g s}^{-1}, \quad \text{in the case of a NS.}$$




Contracting White Dwarf!!!!

A new idea




Contraction of a white dwarf



$M=1.28 \text{ Msolar}$



Comparison with the code of white dwarf evolution by Bergeron et al.







The code by Blinnikov and Dunina-Barkovskaya (1994).

We compared it with recent codes for WD evolution.


Correspondence is very good.



Comparison with BaSTI




Evolution of the moment of inertia



$$I = \frac{8\pi}{3} \int_0^R \rho r^4 dr,$$

Period variation: theory and observations



$$I_1/I_2 = P_1/P_2 \quad \dot{P} = \Delta P/\Delta t = \frac{P}{\Delta t} \left(\frac{I_1}{I_2} - 1 \right)$$

Contraction vs. accretion

Let us compare torque due to accretion onto a WD with the spin-up due to contraction.

Let the accretion disc be terminated at the corotation radius:

$$R_{\text{co}} = (GMP^2/4\pi^2)^{1/3}$$

Then we have

$$2\pi I\dot{\nu} = \dot{M}(GMR_{\text{co}})^{1/2}$$

With accretion rate

$$\dot{M} = L(GM/R)^{-1} = 1.8 \times 10^{14} L_{32} \text{ g s}^{-1}$$

we obtain

$$\dot{\nu} = 1.1 \times 10^{-19} L_{32} I_{50}^{-1} (R/3000 \text{ km})^{-1} \text{ Hz s}^{-1}$$

i.e.

$$\dot{P} = 2 \times 10^{-17} L_{32} \text{ s s}^{-1}$$

WD properties

To fit all data the WD might have:


$M=1.28 M_{\text{solar}}$

Age ~ 2 Myrs ($1 < \text{Age} < 5$ Myrs)


This corresponds to

- radius 3340 km
- temperature 75 000K

Luminosity of the sdO star is much larger $\sim 10^4 L_{\text{solar}}$.




Binary and its evolution



Time (Myr)	$M_1 (M_\odot)$	$M_2 (M_\odot)$	Period (days)	Stage
0.0	7.0	6.75	4550.3	ZAMS
48.8	7.06	6.75	4550.3	RG+MS
49.0	7.05	6.75	4551.6	CHB+MS
53.0	6.89	6.75	4621.7	CHB+RG
53.1	6.89	6.75	4623.4	CHB+CHB
55.0	6.84	6.69	4691.9	EAGB+CHB
55.3	6.8	6.69	4657.4	TPAGB+CHB
55.7	5.96	6.84	4101.8	CE
55.7	1.28	1.47	1.48	ONe WD+He*
64.1	1.28	1.43	1.52	ONe WD+HeG
64.8	1.28	1.42	1.53	CE
64.8	1.28	0.83	0.15	ONe+CO WDs
467.5	1.28	0.83	0.0004	Merger


He-star is formed after the WD


Binary statistics



~25 systems in the Galaxy very similar to HD 49798
and hundreds of systems with slightly different masses.

Pdot for different parameters of WDs





Conclusions

We conclude that all properties of the binary system HD 49798 can be explained in the model where the accreting compact object is a young WD with an age ~ 2 Myrs.

Observed luminosity is due to accretion but observed \dot{P} is due to WD contraction.

The system is rare but not unique. We expect that tens or even hundreds of such binaries are there in the Galaxy.

Discovery of similar system might help to probe interesting early stages of WD evolution.

MNRAS (2018, in press) arXiv: 1711.02449